

Poster presentation

Atazanavir is safe and efficacious in HBV and HCV co-infected patients: results of AI424138 (CASTLE)

J Absalon^{*1}, G Thal², A Thiry¹, R Yang¹, MD Mancini¹ and D McGrath¹Address: ¹Bristol-Myers Squibb Company, Wallingford, CT, USA and ²Bristol-Myers Squibb Company, Lawrenceville, NJ, USA

* Corresponding author

from Ninth International Congress on Drug Therapy in HIV Infection
Glasgow, UK. 9–13 November 2008

Published: 10 November 2008

Journal of the International AIDS Society 2008, **11**(Suppl 1):P136 doi:10.1186/1758-2652-11-S1-P136This abstract is available from: <http://www.jiasociety.org/content/11/S1/P136>

© 2008 Absalon et al; licensee BioMed Central Ltd.

Background

Chronic HBV and HCV infections are common co-morbidities that can complicate antiretroviral treatment in HIV-infected patients. Information on efficacy and safety are necessary to better define optimal therapeutic options in this population.

Methods

Randomized, open-label prospective study comparing once-daily ATV/r with twice-daily LPV/r, both in combination with once-daily fixed dose combination tenofovir/emtricitabine in antiretroviral-naïve HIV-1 infected subjects. Proportion of subjects with HIV RNA <50 c/mL (confirmed virologic response/CVR), changes in CD4 cell count and lipids from baseline, and adverse events (AEs) through 48 weeks are presented among chronic HBV- and/or HCV-infected (Hep+) subjects.

Summary of results

At baseline, 13% of subjects were Hep+ (5% HBV+; 8% HCV+). Overall 9% of females, 14% males, 25% Asians, 15% Blacks, 5% others, and 13% of Whites were Hep+ (Table 1.)

Grade 2–4 treatment related hyperbilirubinemia (15% vs. 0) and jaundice (3% vs. 0) were more common on ATV/r. Nausea (8% vs. 2%) and diarrhea (14% vs. 0) were more common on LPV/r. Few SAEs were reported among Hep+ in either treatment arm. Grade 3–4 elevations in liver function tests were reported among the following: 5/60, 8% (ALT), 5/60, 8% (AST) and 23/60, 38% (total

bilirubin) in the ATV/r arm and 3/50, 6% (ALT), 0% (AST); 0% (total bilirubin) in the LPV/r arm. (Table 2.)

Conclusion

Virologic and immunologic response was comparable in Hep+ treated with ATV/r or LPV/r. ATV/r had a more favorable lipid profile (TC, non-HDL, LDL, TG) and fewer gastrointestinal adverse events among Hep+ subjects than LPV/r. While the overall rates of transaminitis in Hep+ were low in this study compared to those observed in other clinical trials, a small number of subjects in the ATV/r and none on LPV/r had grade 3–4 AST elevations. The cause of this higher proportion in the ATV/r treatment arm among this limited number of subjects is unclear. With close monitoring of liver function tests, ATV/r can be considered as part of HAART among treatment-naïve Hep+ patients.

Table 1:

Efficacy at Week 48	ATV/r N = 61	LPV/r N = 51
HIV RNA <50 c/mL, CVR (Non-Completer = Failure), n/N (%)	42/61 (69)	37/51 (73)
Mean CD4 Cell Count Change from Baseline (SE), cells/mm ³	196 (26.1)	228 (21.7)

**Table 2: Lipid Mean % Change (+/- SE) from Baseline at Week 48
– As Treated Subjects**

	ATV/r N = 60	LPV/r N= 51
Total Cholesterol (TC)	12 (9.4, 15.3)	23 (20.1, 26.4)
HDL	28 (24.2, 32.4)	42 (32.3, 53.1)
Non-HDL	7 (3.1, 10.9)	17 (14.2, 20.9)
LDL	12 (6.4, 16.9)	21 (14.6, 27.8)
Triglycerides (TG)	18 (9.8, 27.1)	45 (33.2, 58.2)

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

